

## Diagnostic Test Solutions

### Problem 1

Consider the following pseudocode:

```
ROUTINE( $n$ )  
1  if  $n = 1$   
2    then return 1  
3    else return  $n + \text{ROUTINE}(n - 1)$ 
```

(a) Give a one-sentence description of what  $\text{ROUTINE}(n)$  does. (Remember, don't guess.)

**Solution:** The routine gives the sum from 1 to  $n$ .

(b) Give a precondition for the routine to work correctly.

**Solution:** The value  $n$  must be greater than 0; otherwise, the routine loops forever.

(c) Give a one-sentence description of a faster implementation of the same routine.

**Solution:** Return the value  $n(n + 1)/2$ .

### Problem 2

Give a short (1–2-sentence) description of each of the following data structures:

(a) FIFO queue

**Solution:** A dynamic set where the element removed is always the one that has been in the set for the longest time.

(b) Priority queue

**Solution:** A dynamic set where each element has an associated priority value. The element removed is the element with the highest (or lowest) priority.

(c) Hash table

**Solution:** A dynamic set where the location of an element is computed using a function of the element's key.

### Problem 3

Using  $\Theta$ -notation, describe the worst-case running time of the best algorithm that you know for each of the following:

(a) Finding an element in a sorted array.

**Solution:**  $\Theta(\log n)$

(b) Finding an element in a sorted linked-list.

**Solution:**  $\Theta(n)$

(c) Inserting an element in a sorted array, once the position is found.

**Solution:**  $\Theta(n)$

(d) Inserting an element in a sorted linked-list, once the position is found.

**Solution:**  $\Theta(1)$

### Problem 4

Describe an algorithm that locates the first occurrence of the largest element in a finite list of integers, where the integers are not necessarily distinct. What is the worst-case running time of your algorithm?

**Solution:** Idea is as follows: go through list, keeping track of the largest element found so far and its index. Update whenever necessary. Running time is  $\Theta(n)$ .

### Problem 5

How does the height  $h$  of a balanced binary search tree relate to the number of nodes  $n$  in the tree?

**Solution:**  $h = O(\lg n)$

**Problem 6**

Does an undirected graph with 5 vertices, each of degree 3, exist? If so, draw such a graph. If not, explain why no such graph exists.

**Solution:** No such graph exists by the Handshaking Lemma. Every edge adds 2 to the sum of the degrees. Consequently, the sum of the degrees must be even.

**Problem 7**

It is known that if a solution to Problem A exists, then a solution to Problem B exists also.

(a) Professor Goldbach has just produced a 1,000-page proof that Problem A is unsolvable. If his proof turns out to be valid, can we conclude that Problem B is also unsolvable? Answer yes or no (or don't know).

**Solution:** No

(b) Professor Wiles has just produced a 10,000-page proof that Problem B is unsolvable. If the proof turns out to be valid, can we conclude that problem A is unsolvable as well? Answer yes or no (or don't know).

**Solution:** Yes

**Problem 8**

Consider the following statement:

If 5 points are placed anywhere on or inside a unit square, then there must exist two that are no more than  $\sqrt{2}/2$  units apart.

Here are two attempts to prove this statement.

**Proof (a):** Place 4 of the points on the vertices of the square; that way they are maximally separated from one another. The 5th point must then lie within  $\sqrt{2}/2$  units of one of the other points, since the furthest from the corners it can be is the center, which is exactly  $\sqrt{2}/2$  units from each of the four corners.

**Proof (b):** Partition the square into 4 squares, each with a side of  $1/2$  unit. If any two points are on or inside one of these smaller squares, the distance between these two points will be at most  $\sqrt{2}/2$  units. Since there are 5 points and only 4 squares, at least two points must fall on or inside one of the smaller squares, giving a set of points that are no more than  $\sqrt{2}/2$  apart.

Which of the proofs are correct: (a), (b), both, or neither (or don't know)?

**Solution:** (b) only

### Problem 9

Give an inductive proof of the following statement:

For every natural number  $n > 3$ , we have  $n! > 2^n$ .

**Solution:** Base case: True for  $n = 4$ .

Inductive step: Assume  $n! > 2^n$ . Then, multiplying both sides by  $(n + 1)$ , we get  $(n + 1)n! > (n + 1)2^n > 2 * 2^n = 2^{n+1}$ .

### Problem 10

We want to line up 6 out of 10 children. Which of the following expresses the number of possible line-ups? (Circle the right answer.)

- (a)  $10!/6!$
- (b)  $10!/4!$
- (c)  $\binom{10}{6}$
- (d)  $\binom{10}{4} \cdot 6!$
- (e) None of the above
- (f) Don't know

**Solution:** (b), (d) are both correct

### Problem 11

A deck of 52 cards is shuffled thoroughly. What is the probability that the 4 aces are all next to each other? (Circle the right answer.)

- (a)  $4!49!/52!$
- (b)  $1/52!$
- (c)  $4!/52!$
- (d)  $4!48!/52!$
- (e) None of the above
- (f) Don't know

**Solution:** (a)

### Problem 12

The weather forecaster says that the probability of rain on Saturday is 25% and that the probability of rain on Sunday is 25%. Consider the following statement:

The probability of rain during the weekend is 50%.

Which of the following best describes the validity of this statement?

- (a) If the two events (rain on Sat/rain on Sun) are independent, then we can add up the two probabilities, and the statement is true. Without independence, we can't tell.
- (b) True, whether the two events are independent or not.
- (c) If the events are independent, the statement is false, because the the probability of no rain during the weekend is  $9/16$ . If they are not independent, we can't tell.
- (d) False, no matter what.
- (e) None of the above.
- (f) Don't know.

**Solution:** (c)

### Problem 13

A player throws darts at a target. On each trial, independently of the other trials, he hits the bull's-eye with probability  $1/4$ . How many times should he throw so that his probability is 75% of hitting the bull's-eye at least once?

- (a) 3
- (b) 4
- (c) 5
- (d) 75% can't be achieved.
- (e) Don't know.

**Solution:** (c), assuming that we want the probability to be  $\geq 0.75$ , not necessarily exactly 0.75.

**Problem 14**

Let  $X$  be an indicator random variable. Which of the following statements are true? (Circle all that apply.)

(a)  $\Pr\{X = 0\} = \Pr\{X = 1\} = 1/2$

(b)  $\Pr\{X = 1\} = E[X]$

(c)  $E[X] = E[X^2]$

(d)  $E[X] = (E[X])^2$

**Solution:** (b) and (c) only